

VSOP OBSERVATION OF THE QUASAR PKS 2215+020: A NEW LABORATORY FOR CORE-JET PHYSICS AT $Z = 3.572$ A. P. LOBANOV¹, L. I. GURVITS^{2,3}, S. FREY⁴, R. T. SCHILIZZI^{2,5},
K. I. KELLERMANN⁶, N. KAWAGUCHI⁷, I. I. K. PAULINY-TOTH¹*To appear in The Astrophysical Journal, v.546, n.2*

ABSTRACT

We report results of a VSOP (VLBI Space Observatory Programme) observation of a high redshift quasar PKS 2215+020 ($z = 3.57$). The ~ 1 milliarcsecond resolution image of the quasar reveals a prominent ‘core-jet’ structure on linear scales from $5h^{-1}$ to $300h^{-1}$ pc ($H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$). The brightness temperatures and sizes of bright features identified in the jet are consistent with emission from relativistic shocks dominated by adiabatic energy losses. The jet is powered by the central black hole with estimated mass of $\sim 4 \times 10^9 M_\odot$. Comparisons with VLA and ROSAT observations indicate a possible presence of an extended radio/X-ray halo surrounding 2215+020.

Subject headings: VLBI — galaxies: radio, quasars: radio, quasars: individual:
PKS 2215+020

1. INTRODUCTION

VLBI⁸ observations of high redshift quasars at centimeter wavelengths allow to address two connected topics:

(1) *Frequency-dependent properties of the milliarcsecond structures in quasars.* An image obtained at a received frequency ν_r represents the source structure at a higher emitted frequency $\nu_{em} = \nu_{obs}(1+z)$ thus facilitating a direct comparison between radio structures at a given frequency in high redshift quasars and their low redshift counterparts observed at a higher frequency ν_{obs} .

(2) *Cosmological tests using the milliarcsecond structure of QSOs as a “standard” object.* As shown in several recent studies, the “apparent angular size – redshift” ($\theta - z$, Kellermann 1993, Gurvits 1994, Gurvits et al. 1999) and “proper

motion – redshift” ($\mu - z$, Vermeulen & Cohen 1994, Vermeulen 1996, Kellermann et al. 1999) relation on milliarcsecond scales have a potential to provide meaningful cosmological information. In this respect, sources at extremely high redshifts are of special interest.

In this paper we present a VSOP (VLBI Space Observatory Programme, Hirabayashi et al. 1998) observation of the high-redshift quasar PKS 2215+020. The observations have been conducted as part of a VSOP study of extremely high redshift quasars. The complete results of this project will be presented elsewhere (Gurvits et al. 2000).

The VSOP is a Space VLBI (SVLBI) mission utilizing an orbiting 8-meter antenna deployed on the Japanese satellite HALCA⁹ (Hirabayashi et al. 1998) and the worldwide array of radio tele-

¹Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany; alobanov@mpifr-bonn.mpg.de, ipauliny@mpifr-bonn.mpg.de

²Joint Institute for VLBI in Europe, P.O.Box 2, 7990 AA, Dwingeloo, The Netherlands; lgurvits@jive.nl, schilizzi@jive.nl

³Astro Space Center of P.N.Lebedev Physical Institute, Moscow, Russia

⁴FÖMI Satellite Geodetic Observatory, P.O. Box 546, H-1373 Budapest, Hungary; frey@sgo.fomi.hu

⁵Leiden Observatory, P.O. Box 9513, 2300 RA, Leiden, The Netherlands

⁶National Radio Astronomy Observatory, 520 Edgemont Rd, Charlottesville, VA 22903-2475, USA; kkellerm@nrao.edu

⁷NAO, Mitaka, Tokyo 181-8588, Japan, kawagu@hotaka.mtk.nao.ac.jp

⁸Very Long Baseline Interferometry

⁹The Highly Advanced Laboratory for Communication and Astronomy

scopes. The satellite has an elliptical orbit, with the apogee and perigee heights at $\simeq 21\,000$ km and $\simeq 560$ km, respectively, and orbiting period of roughly 6 hours. In each observation, data from the satellite are recorded by a network of up to 5 tracking stations, and subsequently correlated with the data from participating ground radio telescopes. Regular VSOP observations started in September 1997 at 1.6 and 5 GHz. A 1.6 GHz VSOP observation of PKS 2215+020 described here was made on September 14-15, 1997.

PKS 2215+020 is an optically faint ($m_B = 21.97$), radio-loud quasar ($S_{5\text{ GHz}} = 0.5\text{--}0.6$ Jy; Griffith & White 1993, Gregory et al. 1996) included in the Parkes half-jansky flat-spectrum sample (Drinkwater et al. 1997). It has an emission redshift, $z = 3.572$ (Drinkwater et al. 1997); a corresponding linear scale of $3.38 h^{-1} \text{ pc mas}^{-1}$ (for the Hubble constant $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ and deceleration parameter $q_0 = 0.5$). The radio spectral index of 2215+020 is $\alpha_{2.7\text{ GHz}}^{5\text{ GHz}} = -0.15$ ($S_\nu \propto \nu^\alpha$). Siebert & Brinkmann (1998) have observed X-ray emission from 2215+020 with the ROSAT High Resolution Imager (HRI), and found only marginal evidence for possible elongation along the P.A. = $60\text{--}70^\circ$.

2. VSOP OBSERVATION AND DATA REDUCTION

We observed the quasar PKS 2215+020 at 1.6 GHz, using the HALCA satellite and a global VLBI array of 15 ground-based radio telescopes (Table 1). The observation lasted for 10 hours, with HALCA data recorded for 7 hours. The data were recorded in the VLBA format (Rogers 1995), with a total observing bandwidth of 32 MHz divided in two intermediate frequency (IF) bands, each having 256 spectral channels. The tracking stations in Green Bank and Goldstone (USA) were used for the HALCA data acquisition. The data were correlated at the VLBA correlator in Socorro (Benson 1995), with a pre-averaging time of 1.966 and 0.524 sec for the ground-ground and space-ground baselines respectively. Fringe visibilities were detected in the HALCA data recorded at both tracking stations. The resulting sampling function (uv -coverage) of the final correlated dataset is shown in Figure 1, indicating an improvement of resolution by a factor of ~ 2.5 compared to ground-based VLBI observations at the same frequency, owing to the ground-space baselines.

We used AIPS¹⁰ and the DIFMAP program (Shepherd et al. 1994) for post-processing of the correlated dataset. The a priori amplitude calibration was determined from the antenna gain and system temperature measurements, including that of the HALCA radio telescope (Moellenbrock et al. 2000). After inspecting the IF bandpasses, the last 46 channels in each IF were deleted, owing to a cutoff in the filter bandpass. This has reduced the effective observing bandwidth to 26.2 MHz. We corrected the residual delays and rates, using single-band (SB) and multi-band (MB) fringe fitting (Cotton 1995) with solution intervals of 5 minutes, and accepted all solutions with $\text{SNR} \geq 7$. To improve the SNR on the space-ground baseline solutions, we fringe fitted the HALCA data, using the source model obtained by imaging the ground-ground VLBI data. After the fringe fitting, the residual phase variations on the ground-ground baselines were within 3° , and within 15° , on the space-ground baselines. We then averaged over all frequency channels, and calibrated the phases with a point source model (to enable time averaging). Finally, the data were exported into DIFMAP and further time-averaged into 60-second bins. The amplitude and phase errors were re-calculated in DIFMAP, from the scatter in the unaveraged data. The estimated RMS noise on the HALCA baselines was about 4 times higher than on the ground-ground baselines, consistent with the smaller antenna size and higher system temperature of the HALCA radio telescope. The amplitude visibility distribution obtained after the amplitude calibration and fringe-fitting is shown in Figure 2.

3. GROUND VLBI ARRAY AND VSOP IMAGES OF 2215+020

From the final fringe-fitted dataset, we produced two images: a) an image with all data included (VSOP image, hereafter), and b) an image using only the ground-ground baseline data included (GVLBI image, hereafter). In both cases, uniform weighting is used, and gridding weights are scaled inversely with the amplitude errors. The resulting images are shown in Figure 3, and their basic characteristics are presented in Table 2.

The images in Figure 3 show a clear core-jet type morphology. We note that

¹⁰The NRAO Astronomical Image Processing System

the general milliarcsecond-scale morphology of PKS 2215+020 described here is consistent with the VLBA 15 GHz image (Kellermann et al. 2000). In the GVLBI image, we identify the core, C, and 7 enhanced emission regions (jet components J1–J7, hereafter) which we will discuss below. The jet extends up to almost 80 mas from the core, and shows a particularly wide, bright section between 45 and 60 mas ($150\text{--}200 h^{-1}\text{pc}$ projected distance), suggesting a possible working surface of a young jet propagating in a dense ambient medium.

The basic structures appear to be similar, in both images shown in Figure 3, taking into account the larger noise and scatter of the data on the space-ground baselines, which results in the increased RMS noise and stronger deconvolution effects in the VSOP image. These effects can be countered by using the natural weighting for the VSOP dataset. We find that the resolution and sensitivity of an image of 2215+020 obtained from the naturally weighted VSOP data are practically identical to those of the uniformly weighted ground VLBI data from our observation (which is commonly expected for VSOP observations; Murphy et al. 1999). Since our goal is to study the jet at highest resolution, we will discuss exclusively the uniformly weighted data represented by the images in Figure 3.

The most interesting difference between the two images shown in Figure 3 is the absence of the component J4 in the VSOP image. The adjacent features J3 and J5 can be identified in both images, even though, in the GVLBI image, they appear to be weaker than J4. It is not immediately clear whether this effect results from unidentified imaging problems, or reflects in fact the true nature of J4 (which may be completely resolved in the VSOP image). Deconvolution effects are visible elsewhere in the two images (for instance, in the VSOP image, J6 appears being split into two features). On the other hand, deconvolution alone is not likely to completely eliminate a feature. In order to provide quantitative criteria for deciding upon this issue, we turn now to modeling the source structure using the interferometric visibility data (Pearson 1995).

3.1. Model fitting the jet structure in 2215+020

To model the jet structure in 2215+020, we first use DIFMAP to fit the ground baseline data by 8 circular Gaussian components (marked in Figure 3). Hereafter, we shall call this model “the GVLBI model”. The GVLBI model is then used as a starting guess for modeling the visibilities in the entire VSOP dataset. The resulting model (“the VSOP model,” hereafter) contains only 7 of the original 8 components. The component corresponding to feature J4 is indeed completely resolved in the VSOP data. In the VSOP model, the formal size of J4 exceeds 20 mas, with the flux of the feature being similar to that in the VLBI model. We take this as evidence in favor of our suggestion that J4 is completely resolved in the VSOP data.

The derived component parameters are listed in Table 3, for both the GVLBI and VSOP models. To estimate the uncertainties of the fitted component parameters, we measure, for each feature, the peak flux density, S_{peak} and its RMS, in the respective image. We then calculate the uncertainties using approximations given by Fomalont (1989).

In several features, both in the GVLBI and VSOP data, the fitted sizes appear to be smaller than the resolution limit. In order to determine which of the fitted values should be considered only as upper limit, we derive, for each feature, its smallest detectable size of a circular Gaussian component in an image with RMS noise σ_{RMS} :

$$d_{\text{lim}} = \frac{\pi}{4} \left[\frac{b_{\text{maj}} b_{\text{min}}}{\ln(2) \ln(S/\sigma_{\text{RMS}})} \right]^{1/2}, \quad (1)$$

where S is the flux density of the feature, and $b_{\text{maj}, \text{min}}$ are the major and minor axes of the restoring beam, respectively. For those features in which $d < d_{\text{lim}}$, we take d_{lim} as the upper limit estimate of the size.

Using the calculated flux densities and sizes, we calculate brightness temperatures, T_{b} , for all jet features.¹¹ For most of the calculated T_{b} , formal errors are of the order of 100% (perhaps indicating that the measured RMS errors of S_{peak} are too conservative). We therefore regard the derived

¹¹We note that our choice of Gaussian components for fitting the visibility data (as opposed to, e.g., optically thick spheroids) does not introduce qualitative changes to the results obtained. Models with non-Gaussian brightness distribution would require only a small, constant correction factor to be applied to the values of brightness temperature presented in this paper.

values of T_b as first order estimates. The component sizes and calculated brightness temperatures are plotted in Figure 4.

4. DISCUSSION

The VSOP observation shows several very interesting properties of the jet in 2215+020. The observed extent of the jet (> 80 mas, $250 h^{-1}$ pc) is remarkable. The structures observed in 2215+020 are almost ten times larger than in any other quasars at $z > 3$ observed with VLBI (c.f. Paragi et al. 1999 and references therein). The morphology of the source suggests that the component J1 may be a working surface of the jet. From the jet expansion presented in Figure 4, we estimate that the jet opening angle should vary between 2° and 7° (with the smaller value obtained not taking into account the two most peculiar features: J4, and J1). These values must also be corrected by a deprojection factor, $\sin \theta_j^{-1}$. However, considering the unusually large extent of the jet, the effect of deprojection is not likely to be strong.

4.1. Brightness temperature

The observed brightness temperatures plotted in Figure 4 can be used to probe the physics of the jet. Following Marscher (1990), we assume that each of the jet components is an independent plane shock in which the radio emission is dominated by adiabatic energy losses. The jet plasma has a power law energy distribution, $N(E)dE \propto E^{-s}dE$, and the magnetic field is assumed to vary as $B \propto d^{-a}$. The Doppler factor is assumed to be constant or varying weakly throughout the jet. Under these assumptions, one can relate the brightness temperature, $T_{b,J}$, of each jet component to the brightness temperature of the core, $T_{b,C}$:

$$T_{b,J} = T_{b,C} (d_J/d_C)^{-\epsilon}, \quad (2)$$

where d denotes the respective measured sizes of the core and jet feature, and $\epsilon = [2(2s + 1) + 3a(s + 1)]/6$. We take $s = 2.0$ (corresponding to synchrotron emission with a spectral index $\alpha = -0.5$) and $a = 1$ (which corresponds to the transverse orientation of magnetic field in the jet). The measured value, $T_{b,C} = 9 \times 10^{11}$, is used for the core. We use the sizes and upper limits measured in the VSOP image, and add the GVLBI measurement for J4. The resulting brightness temperatures are indicated in Figure 4

by thick dashed line. The measured and model values of brightness temperatures agree rather well, suggesting that the jet components may indeed be a collection of relativistic shocks. The discrepancy between the model and observed brightness temperatures is significant only in the outermost jet feature, J1, for which we find a ratio $\xi \equiv T_b^{\text{obs}}/T_b^{\text{model}} \approx 20$. This discrepancy suggests that J1 is a region of the jet where physical conditions may have become substantially different. If J1 interacts with the ambient medium, its material may be going through additional compression and reacceleration, which could lead to a localized increase of the brightness temperature. The observed discrepancy between our model and the measured brightness temperature of J1 may also be explained by variations of the jet Doppler factor, resulting from changes of the speed or orientation of the jet. For this case, one can derive for the Doppler factor of J1, $\delta_{J1} = \delta_j \xi^{1/(\alpha-3)}$ (here δ_j is the Doppler factor in the rest of jet). This condition implies a jet bulk Lorentz factor $\Gamma_j \geq \xi^{1/(3-\alpha)} = 2.5$ and viewing angle $\theta_{\text{jet}} \geq \arccos[(\Gamma_j - \xi^{1/(3-\alpha)})(\Gamma_j^2 - 1)^{-1/2}]$. We find then that the Lorentz factor in J1 must be 40–50% lower than in the rest of the jet (Γ_{J1} is lower than Γ_j because $\delta_j < 1$, so the jet has to be Doppler-deboosted, and its viewing angle must be $\geq 40^\circ$, even for Γ_j as high as 15). Alternatively, if the bulk Lorentz factor is constant throughout the entire jet, the direction of the flow entering the region of J1 must change by 20–30° with respect to the line of sight.

4.2. The transverse size of the jet

For the feature J4, which is resolved out in the VSOP image, the linear size derived from the GVLBI data is ~ 3.4 mas ($12 h^{-1}$ pc). If this size is related to the physical transverse dimension of the jet, the mass of the central object can be estimated, assuming that the jet is collimated by the ambient magnetic field of the host galaxy (e.g. Appl & Camenzind 1992, 1993, Beskin & Malyshev 2000). On parsec scales, the width of a jet can often be underestimated, owing to limited resolution and dynamic range of VLBI observations. In 2215+020, the jet feature J4 is arguably the best measure of the width of the jet, as this feature is the only one resolved out in the VSOP image, and its size (6.2 mas) measured in the GVLBI image is significantly larger

than the respective limiting resolution (2.3 mas). The measured sizes of other jet features (except J1) are most likely affected by the limited resolution and dynamic range of the data, and therefore are likely to be smaller than the true jet width. Finally, J1 is unsuitable for estimating the central black hole mass because it is most likely a strongly perturbed region, actively interacting with the external medium.

For a jet collimated by the ambient magnetic field, B_{ext} , of the host galaxy, the mass of the central object, M_{BH} , can be related to the width of the jet, r_j , so that (Beskin 1997):

$$\begin{aligned} M_{\text{BH}} &\sim 0.5 r_j c^2 G^{-1} (B_{\text{ext}}/B_g)^{1/2} \approx \\ &\approx r_{j,\text{pc}} h^{-1} (B_{\text{ext}}/B_g)^{1/2} 10^{13} M_\odot. \end{aligned} \quad (3)$$

where G is the gravitational constant, R_g is the Schwarzschild radius of the central black hole, and B_g is the magnetic field measured at R_g . Equation (3) refers to the transverse dimension of the jet measured at distances comparable to the collimation scale (typically expected to be located at 100–1000 R_g).

A typical galactic magnetic field is $B_{\text{ext}} \sim 10^{-5}$ G (Beck 2000), and one can expect to have $B_g \sim 10^4$ G (Field & Rogers 1993). With these estimates, we obtain $M_{\text{BH}} \sim 4 \times 10^9 h^{-1} M_\odot$. Since J4 is located at about 80 pc distance from the jet core, its size is most likely affected by the expansion of the jet. This effect may result in overestimating the mass of the central black hole by $\lesssim 30\%$ (assuming a jet opening angle of 4°).¹²

4.3. Large scale radio and X-ray structures in 2215+020

The jet direction observed in our VLBI images is consistent with the extension seen in a VLA observation (Figure 5) on a scale of 7 arcseconds ($23h^{-1}$ kpc). Most interestingly, a similar extension is suggested by the ROSAT HRI image (ROSAT archive directory 701900h¹³). Siebert & Brinkmann (1998) argued that the extension of the emission in the ROSAT image may result from the asymmetrical point-spread function of the HRI. In Figure 6, we overlay the ROSAT and VLA images of 2215+020. Given the remarkable morphological similarity of the X-ray and radio

emission, we tend to conclude that the extension seen in the ROSAT may be real. This would imply that 2215+020 contains an X-ray emitting gas extending out to about $50h^{-1}$ kpc. The X-ray emission may then arise from interaction between the relativistic outflow and a dense circumgalactic gas. It is however not possible to say at present whether the emission on this scales is directly related to the outflow observed on parsec scales. Recent observations with Chandra X-Ray satellite (Marshall et al. 2000) have indicated similar coincidence of radio and X-ray emission distributions in several extragalactic radio sources, most notably in NGC 1275, allowing to speculate that radio lobes may be pushing the X-ray emitting gas outwards. A similar scenario may be at work in 2215+020.

5. SUMMARY

The main results from the observations reported in this paper can be summarized as follows:

1. The radio source PKS 2215+020 has a rich core-jet morphology and unusually large size of the jet which can be traced up to $300h^{-1}$ parsecs. This is by far the longest jet observed at redshifts above 3.

2. The high resolving power of the VSOP allows us to measure the transverse width of the jet, which provides a rough estimate for the mass of the central black hole in the object $M_{\text{BH}} \sim 4 \times 10^9 h^{-1} M_\odot$.

3. The sizes and brightness temperatures derived from the model of the source structure are consistent with the adiabatic energy losses in relativistic shocks embedded in the jet.

4. The arcsecond-scale structure detected in PKS 2215+020 with the VLA coincides with the extension observed in the ROSAT X-ray image, implying a possible presence of an X-ray emitting gas extending out to about $50h^{-1}$ kiloparsecs.

The findings summarized above allow us to present the quasar PKS 2215+020 as a promising laboratory for future studies of core-jet physics at very high redshifts.

We gratefully acknowledge the VSOP Project, which is led by the Institute of Space and

¹²Similar calculations yield $\sim 6 \times 10^8 M_\odot$ for the mass of the central black hole in M87, based on images from ground VLBI observations made at 1.6 GHz (Reid et al. 1989). This estimate is comparable with the value of $\sim 3 \times 10^9 M_\odot$ deduced from HST observations of M87 (Harms et al. 1994).

¹³<ftp://ftp.xray.mpe.mpg.de/rosat/archive/700000/701900h/>

Astronautical Science (Japan) in cooperation with many agencies, institutes, and observatories around the world. We thank W. Brinkmann and M. Gliozzi for their helpful remarks on the interpretation of the ROSAT data. We would like to thank the anonymous referee for comments and suggestions for improving the manuscript. SF acknowledges support from the Hungarian Space Office, NWO and OTKA. LIG acknowl-

edges partial support from the European Commission TMR Programme, Access to Large-Scale Facilities under contract ERBFMGEXT950012, the European Commission, TMR Programme, Research Network Contract ERBFMRXCT 96-0034 "CERES". The National Radio Astronomy Observatory is operated by Associated Universities, Inc. under a Cooperative Agreement with the National Science Foundation.

REFERENCES

- Appl, S. & Camenzind, M. 1992, *A&A*, 256, 354.
 Appl, S. & Camenzind, M. 1993, *A&A*, 274, 699.
 Beck, R. 2000, in "Perspectives on Radio Astronomy", eds. M.P. van Haarlem & J.M. van der Hulst, in press.
 Benson, J.M. 1995, p. 117 in ZDN95.
 Beskin, V.S. 1997, *Physics-Uspekhi*, 40(7), 659.
 Beskin, V.S. & Malyshev, L.M. 2000, *Astronomy Letters*, 26, 208.
 Cotton, W.D. 1995, p. 189 in ZDN95.
 Drinkwater, M.J., Webster, R.L., Francis, P.J., et al. 1997, *MNRAS*, 284, 85.
 Field, G.B. & Rogers, A.D. 1993, *ApJ*, 403, 94.
 Fomalont, E. 1989, in R.A. Perley, F.R. Schwab, & A.H. Bridle (eds.) "Synthesis imaging in radio astronomy", ASP Conference Series, vol. 6, p. 213.
 Gregory, P.S., Scott, W.K., Douglas, K., & Condon, J.J. 1996, *ApJS*, 103, 427.
 Griffith, M.R. & White, A.E. 1993, *AJ*, 105, 1666.
 Gurvits L.I. 1994, *ApJ* 425, 442
 Gurvits L.I., Kellermann K.I., & Frey S. 1999, *A&A*, 342, 378.
 Gurvits L.I., et al. 2000, in preparation.
 Hirabayashi H., Hirose, H., Kobayashi, H., et al. 1998, *Science*, 281, 1825.
 Kellermann K.I. 1993, *Nature* 361, 134
 Kellermann, K.I., Vermeulen, R.C., Zensus, J.A., Cohen, M.H., & West, A. 1999, *New Astronomy Reviews*, 43, 757.
 Kellermann, K.I. et al. 2000, (in preparation).
 Marscher, A.P. 1990, in *Parsec-scale Radio Jets*, eds. J.A. Zensus, & T.J. Pearson (Cambridge: Cambridge University Press), p. 236.
 Marshall, H.L., Canizares, C.R., & Fang, T. 2000, in "High-Energy Astrophysical Phenomena Revealed by Space-VLBI", eds. H. Hirabayashi, P.G. Edwards, & D.W. Murphy (Sagamihara, Japan: ISAS), p. 207.
 Moellenbrock, G.A., Kobayashi, H., Murphy, D.W., et al. 2000, *Advances in Space Research*, 26, 613.
 Murphy, D.W., Tingay, S.J., Preston, R.A., et al. 1999, *New Astronomy Reviews*, 43, 727.
 Paragi, Z., Frey, S., Gurvits, L.I., et al. 1999, *A&A*, 344, 51.
 Pearson, T.J. 1995, p. 309 in ZDN95.
 Reid, M.J., Biretta, J.A., Junor, W., et al. 1989, *ApJ*, 336, 112.
 Shepherd M. C., Pearson, T. J., & Taylor, G. B. 1994, *BAAS* 26, 987.
 Siebert, J. & Brinkmann, W. 1998, *A&A*, 333, 63.
 Rogers, A.E.E. 1995, p. 93 in ZDN95.
 Vermeulen R.C., Cohen M.H. 1994, *ApJ* 430, 467
 Vermeulen R.C. 1996, in *Proc. IAU Symp.* 175, ed. Ekers R. et al., Dordrecht: Kluwer Acad. Publ., p57
 ZDN95: Zensus, J.A., Diamond, P.J., & Napier, P.J. (eds.) 1995, "Very Long Baseline Interferometry and the VLBA", ASP Conference Series, Vol. 82.

FIG. 1.— The (u, v) -coverage of the observation of 2215+020. The largely horizontal, inner tracks are from the ground-ground baselines; the inclined, outer sections are from the ground-space baselines.

FIG. 2.— The distribution of the correlated flux versus uv -distance. The ground-ground baselines extend up to $50 \text{ M}\lambda$; the ground-space baselines cover the uv -distances between 20 and $150 \text{ M}\lambda$.

FIG. 3.— The GVLBI (top) and VSOP (bottom) images of 2215+020. Image characteristics are summarized in Table 2. The lowest positive contours represent the brightnesses of 0.7 mJy/beam (GVLBI image) and 1.2 mJy/beam (VSOP image). Contour levels are drawn at $-1.0, 1.0, 1.41, 2.0, 2.82, 4, \dots$ of the lowest positive contours. Shown in both images are locations of the Gaussian components obtained from model fitting the structures in the GVLBI image. The model fits are described in section 3.1. In the VSOP image: the component J4 is resolved out; the component J6 appears to be double, likely due to deconvolution errors owing to the much larger noise on the ground-space baselines.

FIG. 4.— Sizes and brightness temperatures of different emitting components in the jet of 2215+020. The component identification is described in Figure 3. Open symbols denote upper limits of the size and lower limits of the brightness temperature. The feature J4 is completely resolved in the VSOP image. In the brightness temperature plot, the thick dashed line represents values predicted for relativistic shocks in which adiabatic energy losses dominate the radio emission.

FIG. 5.— A VLA A-array image of 2215+020 at 5.0 GHz , made from an observation on February 26 1993. The restoring beam (2.84×2.64 arcseconds at $\text{P.A.} = 28^\circ$) is plotted in the lower left corner. Contour levels are drawn at $-0.03, 0.03, 0.06, \dots, 61.4\%$ of the peak brightness of 0.606 Jy/beam .

FIG. 6.— ROSAT HRI image of 2215+020 (contours) overlaid on the VLA image (grey scale). The contour levels are drawn at 10, 20, 40, and 80% of the peak of the X-ray emission.

TABLE 1

SELECTED TECHNICAL PARAMETERS OF RADIO TELESCOPES PARTICIPATED IN THE VSOP OBSERVATION OF 2215+020.

Telescope	Code ¹	D ² [m]	T _{sys} ³ [K]	K ⁴ [K/Jy]	SEFD ⁵ [Jy]
HALCA Satellite		8	75	0.0043	17400
Effelsberg, Germany	Eb	100	40	1.50	30
Medicina, Italy	Mc	32	70	0.10	700
Noto, Italy	Nt	32	95	0.11	950
Torun, Poland	Tr	32	50	0.13	380
Green Bank, USA	Gb	43	30	0.29	90
VLBA, USA:					
Saint Croix	Sc	25	28	0.082	310
Hancock	Hn	25	33	0.080	330
North Liberty	Nl	25	20	0.071	290
Fort Davis	Fd	25	24	0.090	270
Los Alamos	La	25	32	0.121	280
Pietown	Pt	25	27	0.101	270
Kitt Peak	Kp	25	31	0.099	310
Owens Valley	Ov	25	34	0.100	340
Brewster	Br	25	24	0.087	280
Mauna Kea	Mk	25	34	0.097	340

NOTE.—1 – two-letter station code; 2 – antenna diameter; 3 – receiver system temperature; 4 – antenna sensitivity; 5 – system equivalent flux density.

TABLE 2

CHARACTERISTICS OF THE GROUND ARRAY AND VSOP IMAGES

	Image:	
	GVLBI	VSOP
Beam size [mas]	9.5×2.8	4.5×1.1
Total CLEAN flux [mJy]	451.3	465.8
Map peak [mJy/beam]	277.8	221.4
Map noise [mJy/beam]	0.16	0.36
Thermal noise [mJy/beam]	0.08	0.07

TABLE 3
MODEL FITS OF THE STRUCTURES IN THE GROUND ARRAY AND VSOP IMAGES

(1)	(2)	S_{peak} [mJy/beam] (3)	S [mJy] (4)	r [mas] (5)	ϕ [deg] (6)	d [mas] (7)	d_{lim} [mas] (8)	T_b [K] (9)
C:	GVLBI	278±69	291±100	0.0	...	(0.9)	1.8	$> 1.9 \times 10^{11}$
	VSOP	221±81	300±137	0.0	...	0.8±0.3	0.8	9.1×10^{11}
J7:	GVLBI	4.2±2.3	11.6±6.8	3.3±0.6	88±11	(<0.1)	2.3	$> 4.4 \times 10^9$
	VSOP	3.7±2.0	10.0±5.8	3.0±0.3	93±6	(<0.1)	1.2	$> 1.5 \times 10^{10}$
J6:	GVLBI	2.6±1.6	3.1±2.5	7.8±0.9	88±6	(<0.1)	2.8	$> 8.1 \times 10^8$
	VSOP	2.1±1.1	2.8±1.8	6.9±0.4	84±3	(<0.1)	1.5	$> 2.5 \times 10^9$
J5:	GVLBI	2.5±1.3	3.9±2.4	14.6±0.4	90±3	(1.9)	2.7	$> 1.1 \times 10^9$
	VSOP	2.5±1.2	3.4±2.0	12.0±0.4	89±2	1.7±0.8	1.4	2.0×10^9
J4:	GVLBI	2.8±1.4	12.1±6.2	23.7±1.6	78±4	6.2±3.1	2.3	2.1×10^8
	VSOP			<i>resolved</i>				
J3:	GVLBI	1.6±0.8	11.3±5.7	34.0±1.0	91±2	4.0±2.0	2.3	5.0×10^8
	VSOP	1.8±0.8	9.4±4.3	32.1±0.8	85±2	3.8±1.7	1.2	1.4×10^9
J2:	GVLBI	9.1±3.5	12.9±6.1	49.6±0.6	84±1	3.3±1.3	2.3	7.4×10^8
	VSOP	3.8±1.7	12.0±5.6	49.0±0.7	87±1	3.1±1.4	1.1	2.6×10^9
J1:	GVLBI	18.0±5.4	110±34	59.5±2.1	74±2	13.8±4.1	1.9	4.2×10^7
	VSOP	5.7±2.8	123±60	57.1±3.0	77±2	12.2±6.0	0.9	1.6×10^8

NOTE.—Column designation: 3 – component peak brightness (from the images); 4 – component flux density; 5 – distance from the core C; 6 – position angle; 7 – component size (values in brackets denote upper limits; listed sizes of <0.1 mas denote tentative values for the features in which the model fitting has failed to produce a viable estimate); 8 – limiting size (from equation 1); 9 – brightness temperature.

2215+020 at 1.634 GHz 1997 Sep 14











